

OXYTETRA 200 L.A.

INJECTABLE SOLUTION

Composition:

Each ml solution contains:

200 mg oxytetracycline base (as dihydrate)

Spectrum of activity:

Oxytetracycline is a broad spectrum antibiotic, with bacteriostatic action against a large number of Gram-positive and Gram-negative organisms, (like streptococci, staphylococci, *Pasteurella* spp., *Brucella* spp., *Corynebacteria* spp., *Actinobacillus* spp., *Eperythrozoon* spp., *Bordetella* spp., *Ehrlichia* spp., coliforms, *Salmonella* spp.) and spirochetes, mycoplasmas, *Rickettsiae* spp., *Chlamydia* spp. and some protozoa. Activity against *Pseudomonas*-, *Proteus*- and *Klebsiella* spp. is low. Applied parenterally, oxytetracycline penetrates quickly into the blood and tissues, but the advanced effect of this preparation is the extremely long therapeutic effect.

Indications:

Infections caused by micro-organisms susceptible to oxytetracycline, like infections of the respiratory tract (pneumonia, bronchopneumonia), infections of the urinary tract, enteritis, (endo)metritis, mastitis, foot rot and Bedsoniasis in cattle, pigs, sheep, goats and poultry.

Contra-indications:

Hypersensitivity to tetracyclines, serious kidney and/or liver disturbances, combination with penicillins or cephalosporins, use between 5 days before till 5 days after vaccination.

Side-effects:

Hypersensitivity reactions, irritation at injection site.

Dosage and administration:

The general dose is: 20 mg/kg bodyweight (= 1 ml per 10 kg bodyweight) by deep i.m. injection. Three days action. Avoid application of more than 10 ml per injection site. In severe cases inj. can be repeated after 3 days. Piglets till 10 kg maximal 1 ml subcutaneous.

Withdrawal time:

For meat: 21 days.

For milk: 7 days.

Storage:

Store between 15 °C and 25 °C and protected from light.

Shelf life:

3 years.

Presentation:

100 ml multidose vial.



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN