

SULFADINE 333

INJECTABLE / ORAL SOLUTION

Composition:

Each ml solution contains:

333 mg sulfadimidine sodium

Spectrum of activity:

Sulfadimidine is a sulphonamide with antimicrobial activity against Gram-positive cocci (haemolytic streptococci, pneumococci and some staphylococci), against some Gram-negative bacteria (like *Pasteurella* spp., *E. coli*, *Salmonella* spp.) and against some *Rickettsiae* and protozoa. Bacterial resistance against sulphonamides mostly is permanent and cross-resistance between sulphonamides is common.

Indications:

Pneumonia, foot rot, diphtheria in calves, actinobacillosis, colibacillosis, salmonellosis, pasteurellosis, enteritis, polyarthritis, streptococcal mastitis, dysentery, lymphangitis, septicaemia in young animals and coccidiosis in horses, cattle, pigs, sheep, goats and poultry. Sulfadimidine is rapidly absorbed from the site of application and slowly excreted by the organism or animals, so in most animals a single daily dose is sufficient.

Contra-indications:

Liver or renal dysfunctions, diseases with decreased urine production (oliguria, anuria), hypersensitivity to sulphonamides. Sulfadimidine is incompatible with para-aminobenzoic acid and its esters (e.g. procaine or tetracaine).

Side-effects:

Sometimes haemorrhagic enteritis (in cattle), kidney damage, hypersensitivity reactions or crystalluria.

Dosage and administration: For slow intravenous, intramuscular or oral administration.

Cattle, sheep, goats, pigs, horses: 3 ml per 10 kg b.w. the first day, followed by 1.5 ml per 10 kg during 3 – 5 days.

Oral administration. Poultry: 5 – 6 ml per litre of drinking water, during 3 – 6 days. Cattle: 3 ml per 10 kg b.w. during 3 – 6 days. Provide ample drinking water during treatment.

Withdrawal time:

For milk: 5 days.

For meat: 8 days (horse, cow, sheep, goat)
10 days (calf, pig, poultry).

Storage:

Store at 15 -25 °C and protected from light.

Shelf life:

2 years (250 – 500 ml)

3 years (100 ml)

Presentation:

100 ml multi dose vial. Plastic vial 250 ml and 500 ml.



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN