

TYLOCARE

INJECTABLE SOLUTION

Composition:

Each ml solution contains:
200 mg tylosin base (as tartrate).

Spectrum of activity:

Tylosin, a macrolide antibiotic, is active against particularly Gram-positive bacteria like *Streptococcus* spp. and *Staphylococcus* spp. (including *Staphylococcus aureus*), some Spirochetes (including *Leptospira*); *Actinomyces*, Mycoplasmas (PLO), *Haemophilus pertussis*, *Moraxella bovis* and some Gram-negative cocci.

Indications:

Infections caused by micro-organisms susceptible to tylosin, like e.g. respiratory tract infections, Dysentery Doyle and arthritis caused by mycoplasma, mastitis and endometritis. After parenteral administration, therapeutically active blood concentrations of tylosin are reached within 2 hours. Target animals: cattle, sheep, pigs, poultry and dogs.

Contra-indications:

Hypersensitivity to tylosin, cross-hypersensitivity to macrolides. Use in laying hens.

Side-effects:

Sometimes, local irritation at the injection site may occur (inject slowly).

Dosage and administration:

For intramuscular or subcutaneous administration.
The general dose is: 0.5 -1 ml per 20 kg bodyweight daily, during 3-5 days.

Withdrawal time:

For meat: 8 days (cattle, sheep).
4 days (pigs, poultry).
For milk: 4 days.

Storage:

Store between 15 °C and 25 °C and protected from light.

Shelf life:

2 years.

Presentation:

100 ml and 250 ml multidose vial.



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN