

TOLTRAPAN

ORAL SOLUTION

Composition:

Each ml solution contains:
25 mg toltrazuril

Spectrum of activity:

Toltrazuril, a triazinetrione derivative, is active against all intracellular stages of coccidia, including schizonts, micro- and macrogamontes.

Toltrazuril acts against the following coccidian of poultry: Chicken: *Eimeria acervulina*, *Eimeria brunette*, *Eimeria Maxima*, *Eimeria mitis*, *Eimeria necatrix*, *Eimeria tenella*. Turkey: *Eimeria adenoids*, *Eimeria meleagridis*, *Eimeria meleagrimitis*

Indications:

Prevention and control of coccidiosis, caused by microorganisms susceptible to toltrazuril in poultry.

Contra-indications:

Usage in laying hens, where eggs are to be used for human consumption.

Side-effects:

No side-effects are reported when used at therapeutic doses.

Dosage and administration:

Oral administration via the drinking water to broilers, layer pullets before 16 weeks age, and turkey flocks.
General dose: 7 mg toltrazuril per kg bodyweight for 2 consecutive days. This generally corresponds with 300 ml TOLTRAPAN per 100 liters of drinking water daily. If necessary repeat treatment after 5 days.

Withdrawal time:

For meat (broilers, turkeys): 21 days.

Storage:

Store closed at 15-25 °C, protected from light.

Presentation:

Plastic bottle of 100 ml and 1 litre.



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN