

AMPROLE 200

WATER SOLUBLE POWDER

Composition:

Each gram powder contains:
200 mg amprolium hydrochloride

Spectrum of activity:

Amprolium is an anti-coccidial drug with activity against *Eimeria tenella*, *E. necatrix*, *E. acervulina*.

The drug acts by interfering with thiamine metabolism in the parasite.

Indications:

As preventive or therapeutic agent against Eimeria infections.

Contra-indications:

Use in hens producing eggs for human food.

Side-effects:

Long-term treatment of high doses may result in delayed growth or polyneuritis (caused by reversible thiamine deficiency) and the development of natural immunity may be delayed.

Dosage and administration:

Orally via drinking water or feed.

Broilers: 15-20 mg/kg b.w. (approx. 60 g per 100 litres of drinking water) during 5-7 days, followed by 7.5-15 mg/kg b.w. (approx. 30 g per 100 litres of drinking water), during 1 or 2 weeks. Administer during treatment only medicated drinking water.

Lambs: 50 mg/kg bodyweight (approx. 5 g powder per 20 kg b.w.), during 1 or 2 days, followed by 5-10 ppm feed (25-50 mg powder/kg feed) during 3 weeks.

Cattle: 50 mg/kg bodyweight (approx. 5 g powder per 20 kg b.w.), during 5 days (via drinking water).

Mixed with feed, the product should be used immediately. Medicated drinking water should be used within 24 hours.

Withdrawal time:

For meat: 2 days.

Storage:

Store closed and dry at 15-25 °C.

Shelf life:

3 years.

Presentation:

Sachet of 100 gram, plastic jar of 1000 gram.



FOR VETERINARY USE ONLY