

# **FLUMESOLE 100**

## **WATER SOLUBLE POWDER**

**Composition:**

Each gram powder contains:  
100 mg flumequine

**Spectrum of activity:**

Flumequine is a quinolone derivative, with bactericidal action against particularly Gramnegative bacteria (e.g. *Klebsiella* spp., *E. coli*, *Enterobacter cloacae*, *Serratia*, *Proteus* spp., *Salmonella* spp., *Neisseria meningitides*, *Pasteurella multocida*, *Campylobacter* spp. and *Shigella* spp.).

**Indications:**

Infections caused by micro-organisms susceptible to flumequine.  
E.g. prevention and treatment of: colibacillosis (*E. coli* infections), pasteurellosis (Cholera) and salmonellosis in poultry, calves and pigs.

**Contra-indications:**

Hypersensitivity to flumequine, serious liver- or renal insufficiency. Use in birds producing eggs for human food.

**Side-effects:**

Eventually hypersensitivity to flumequine may occur.

**Dosage and administration:**

Orally via drinking water. Medicated water should be used within 24 hours.  
Medication should be continuous during the treatment period.

Poultry: 10 mg per kg bodyweight, during 3-5 consecutive days.

This is achieved by adding 100 gram powder per 120 litres of drinking water.

Calves/pigs: 6 mg/kg b.w., twice a day for 5-6 days. This is approximately 500 g per 1000 litres of drinking water for an animal of 50 kg).

**Withdrawal time:**

For meat: 2 days.

**Storage:**

Store closed in a cool and dry place.

**Shelf life:**

3 years.

**Presentation:**

Sachet of 100 gram, plastic jar of 1000 gram.

**FOR VETERINARY USE ONLY**

