

LEVADEX

WATER SOLUBLE POWDER

Composition:

Each gram powder contains:
200 mg levamisole hydrochloride

Spectrum of activity:

Levamisole is a synthetic broad spectrum anthelmintic and is active against adult and larval stages of gastro-intestinal nematodes and lungworms of cattle, sheep, goats and pigs. After oral or parenteral administration, levamisole first paralyzes the worms and afterwards the worms are expelled in the faeces (within 24 hours after parenteral administration). In therapeutic dose levamisole is non-toxic and no side-effects are observed. Levamisole may be given to pregnant animals or animals in poor condition.

Indications:

Adult and larval stages of gastro-intestinal nematodes and lungworms of cattle, sheep, goats, pigs and poultry:

Lungworms (*Dictyoacaulus* spp., *Protostrongylus* spp. and *Metastrongylus elongatus* (pigs)).

Ascaridae (*Toxocara vitulorum*, *Ascaris suum*).

Trichuridae (*Trichuris ovis*, *Trichuris suis*).

Trichostrongylidae (*Trichostrongylus* spp., *Cooperia* spp., *Ostertagia* spp., *Marshallagia* spp., *Haemonchus* spp., *Nematodirus* spp.).

Strongyloidae (*Oesophagostomum* spp., *Chabertia ovina*, *Bunostomum* spp.).

Strongyloididae (*Strongyloides* spp.).

Contra-indications:

Simultaneous administration with substances having nicotinic like action e.g. pyrantel, morantel.

Dosage and administration:

For oral administration with the drinking water.

Cattle, sheep, goats and pigs: the general dose is 10 mg/kg bodyweight (10 gram powder per 200 kg b.w.).

Poultry: 100 ppm. This is achieved by adding 100 gram powder per 200 L of drinking water

As a rule, one single treatment will be sufficient.

Withdrawal time:

For meat: 8 days.

For milk: 3 days.

Storage:

Store at 15-25°C, in a dry place.

Shelf life:

3 years.

Presentation:

Sachet of 100 gram, plastic jar of 1000 gram.

**FOR VETERINARY USE ONLY**