

OXYTETRA 200

WATER SOLUBLE POWDER

Composition:

Each gram powder contains:
200 mg oxytetracycline hydrochloride

Spectrum of activity:

Oxytetracycline is a broad spectrum antibiotic, with bacteriostatic action against a large number of Gram-positive and Gram-negative organisms, (like streptococci, staphylococci, *Pasteurella* spp., *Brucella* spp., *Corynebacteria* spp., *Actinobacillus* spp., *Eperythrozoon* spp., *Bordetella* spp., *Ehrlichia* spp., coliforms, *Salmonella* spp.) and spirochetes, mycoplasmas, *Rickettsiae* spp., *Chlamydia* spp. and some protozoa. Activity against *Pseudomonas*-, *Proteus*- and *Klebsiella* spp. is low.

Indications:

Gastro-intestinal infections (*E. coli*); enteritis (*Salmonella enteritidis*, *Clostridium perfringens*); disturbances of the microbial gutflora (*Streptococcus faecalis*, *Lactobacillus*); atrophic rhinitis (*Pasteurella multocida*, *Bordetella bronchiseptica*); enzootic pneumonia (*Mycoplasma hyopneumonia*) and secondary bacterial infections (*Pasteurella multocida*, *Bordetella bronchiseptica*, *Actinobacillus pleuropneumonia* and *Haemophilus parasuis*).

Contra-indications:

Oral administration to mature animals which have a microbiological digestion in the gastro-intestinal system which is essential (horse, cows, guinea-pigs and hamsters). Administration to animals which are hypersensitive for tetracyclines or to animals with serious renal and liver impairments.

Dosage and administration:

For oral administration via the drinking water.
Calves: 10-20 g per 200 kg bodyweight every 12 hours during 3-5 days.
Pigs: 1-2 g per litre drinking water during 3-5 days.
Poultry (broilers): 1-6 g per litre drinking water during 3-7 days.
Poultry (laying hens): 1-3 g per litre drinking water during 3-5 days.

Withdrawal time:

Meat: Calves: 8 days
Pigs: 5 days.
Poultry: 8 days.
Eggs: 4 days

Storage:

Store closed in a cool and dry place.

Shelf life:

3 years.

Presentation:

Sachet of 100 gram, plastic jar of 1000 gram.



FOR VETERINARY USE ONLY