

PANTACOX

WATER SOLUBLE POWDER

Composition:

Each gram powder contains:

200 mg	sulphamerazine sodium
25 mg	sulphaquinoxaline sodium
25 mg	pyrimethamine
100 mg	furaltadone HCl
5 mg	vitamin K3
15,000 i.u.	vitamin A

Spectrum of activity:

Sulphonamides demonstrate synergy with the dihydrofolate reductase inhibitor pyrimethamine which inhibit a later stage in folic acid synthesis. This combination acts bacteriostatic, and in many cases bactericidal against a large number of Gram-positive and Gram-negative bacteria.

Sulphaquinoxaline is an anticoccidial agent. Furaltadone is active mainly against Gram-negative bacteria, while vitamin K3 and A act as essential nutritional agents.

Indications:

For the prevention and control of caecal and intestinal coccidiosis, Pullorum disease and fowl typhoid enteritis in poultry.

Contra-indications:

Birds producing eggs for human consumption.

Dosage and administration:

Orally, via the drinking water.

The general dose is 1 g/l of drinking water for 3 consecutive days.

Withdrawal time:

For meat: 14 days.

Storage:


Store closed in a cool and dry place.

Shelf life:

3 years.

Presentation:

Sachet of 100 gram, plastic jar of 1000 gram.



FOR VETERINARY USE ONLY