

AMOXY-C

WATER SOLUBLE POWDER

Composition:

Each gram powder contains:

200 mg amoxicillin base (as trihydrate)
1,000,000 I.U. colistin base (as sulphate) (equivalent to 50 mg)

Spectrum of activity:

Amoxicillin is a broad spectrum, semi-synthetic penicillin with activity against both Grampositive and Gramnegative bacteria. Sensitive are streptococci, *E. coli*, *Pasteurella* spp., *Proteus* spp., *Salmonella* spp., *Corynebacterium* spp., staphylococci (β -lactamase negative), *Clostridium* spp.

Colistin, a polymyxin antibiotic, is a small spectrum antibiotic with bactericidal activity against Gramnegative bacteria, e.g. *E. coli*, *Pseudomonas aeruginosa*, *Salmonella* spp., *Haemophilus* spp., *Pasteurella* spp., *Brucella* spp., *Enterobacter aerogenes*, *Shigella* spp. and *Campylobacter* spp. After oral administration, the absorption from the intestines is very poorly.

Indications:

Infections caused by micro-organisms susceptible to amoxicillin and/or colistin in chicken and turkeys, like necrotic enteritis (*Clostridium perfringens*), enteritis and diarrhoea (*Salmonella*, *E. coli*), or infectious coryza (*Haemophilus paragallinarum*). Gastrointestinal infections caused by *E. coli*, *Salmonella*, *Clostridium* and *Campylobacter* spp in pigs, calves and lambs.

Contra-indications:

Hypersensitivity to amoxicillin and/or colistin, renal dysfunctions, combination with fast-acting bacteriostatic antimicrobial drugs (e.g. tetracyclines, chloramphenicol, sulfonamides).

Dosage and administration:

Orally, via drinking water.

Poultry: 1 kg per 4000 litres of drinking water during 3-5 consecutive days.

Pigs/calves/lambs: 5 g per 50 kg body weight during 3-5 consecutive days.

Withdrawal time:

Meat: 6 days (poultry) and 8 days (pigs, calves and lambs).

Storage:

Store closed and dry at 15-25 °C.

Shelf life:

3 years.

Presentation:

Plastic jar of 1000 gram.



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN