

PANTILMI 250

ORAL SOLUTION

Composition:

Each ml solution contains: 250 mg tilmicosin base (as phosphate)

Spectrum of activity:

Tilmicosin is a macrolide antibiotic synthesised from tylosin. It has an antibacterial spectrum similar to tylosin with enhanced activity against *Pasteurella multucida* and *Pasteurella Haemolytica* as well as Mycoplasma spp.

Indications:

For the use in poultry, tilmicosin is indicated for the treatment and control of respiratory diseases associated with mycoplasma and other organisms susceptible to tilmicosin.

Contra-indications:

Usage in laying hens, where eggs are to be used for human consumption.

Side-effects:

No side-effects are reported when used at therapeutic doses.

Dosage and administration:

Oral administration via the drinking water to broilers, layer pullets before 16 weeks age, and turkey flocks.

General dose: 2 ml per 10 L of drinking water daily for 3 days.

Medicated drinking water should be used within 24 hours and prepared fresh every day.

Withdrawal time:

For meat (broilers, turkeys): 10 days.

Storage:

Store closed in a cool place.

Presentation:

Plastic bottle of 100 ml, 500 ml and 1 litre.



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN